

from a continuous-duty, argon-ion laser with two strong lines available, 4880 and 5145 nm. Interaction between the negative ions and the light beam occurs within the laser cavity. The kinetic energy of free electrons photodetached from the negative ions is measured with a hemispherical energy analyzer and an electron multiplier.

The experiment has also been run with hydrogen and deuterium negative ions, whose electron affinities are well known, as references, and the helium electron affinity has been found to be 80.0 ± 2 mV. The absolute total cross section for the photodetachment process was not determined from these experiments. It was found that detachment into the 2^3S state of neutral helium is seven times more likely than detachment into the 2^3P state, for electron ejection essentially parallel to the polarization of the optical field.

Self-Induced Transparency Found in Sulfur Hexafluoride

By applying a short, intense pulse from a Q-switched carbon dioxide laser to gaseous sulfur hexafluoride, Kumar Patel and Richard Slusher of Bell Telephone Laboratories have made the gas transparent to 10.6-micron radiation (*Phys. Rev. Letters* 19, 1019, 1967). They have thus verified the self-induced transparency first found with a ruby laser system by Erwin Hahn and Samuel McCall of Berkeley (*PHYSICS TODAY*, August 1967, page 47). To observe the effect, one must match the exciting frequency to a strongly absorbing transition in the sample.

Unlike the Berkeley experiments, which were plagued by the erratic behavior of pulsed ruby sources, the density of absorbers can be easily varied in the gas to see how transmission varies with collision relaxation times of the absorbing levels, the inhomogeneous Doppler line width and the absorption coefficients.

In Hahn and McCall's theory of the effect, the light beam coherently excites the molecules; they interact with each other and emit light; this process delays and reshapes the pulse into one in which the electric field vector varies

as $\text{sech}(t/\tau)$ (where τ is final pulse width) and under certain conditions widens the pulse. They make the simplification that the pulse is short compared with any relaxation time.

Patel and Slusher verified qualitatively the theory at sulfur hexafluoride pressures from 0.01 to 0.05 torr, and with intensities as low as 10 watts/cm². Then by adding a buffer gas (helium or hydrogen) to the sulfur hexafluoride they shortened the homogeneous relaxation time. They found that the pulse, besides being delayed (to an apparent speed in the sample $1/20$ the speed of light) and reshaped, was narrowed by as much as one half.

In preliminary work Patel and Slusher tried to excite ammonia and ethylene with their carbon dioxide laser but did not succeed because absorption was too low.

The Bell physicists feel that they now have a promising technique for studying relaxation times of individual transitions in gases and gas mixtures.

Somewhere in the universe there may be short, intense, light bursts that pass through gases in space and may produce self-induced transparency in the dilute interstellar medium. Then an astronomer on earth would think the light source (such as the puzzling hydroxyl emitters, (*PHYSICS TODAY*, November 1967, page 69), is a lot farther away than it really is.

Venus Probes Report: Earth is the Oddball

The atmosphere of Venus is almost all carbon dioxide; below the atmosphere the climate is very dry and hot, and the pressure is 15 to 20 times that on earth. This is the forbidding picture of Venusian meteorological conditions that has emerged from the data sent back by Russia's Venus IV spacecraft and by the US Mariner V.

Spectroscopic observations from the ground had already shown the presence of carbon dioxide on Venus, but the concentration was not known. Estimates made since the first observation in 1934 have fluctuated widely between 1 and 90%. Measurements of the thermal infrared emission showed that the temperature is similar to the earth's at the top of the Venusian atmosphere, 235°K, but observations indicated that it is much hotter

at the surface—perhaps 600°K.


Data from the two spacecraft agree that the carbon dioxide concentration is around 90% and that there is no detectable water and apparently only slight traces of oxygen. In addition the Russian Venus probe measured an atmospheric pressure rising from 5 atmospheres at 30 km to about 20 atm at ground level and a temperature range of 270°K at 25 km to 553°K at the surface. This high surface temperature could be an example of the "greenhouse effect;" the Venus atmosphere may be opaque in the infrared but transparent at other frequencies so sunlight is absorbed by the ground and the energy reemitted by blackbody infrared radiation is trapped in the atmosphere. Alternatively it may be due to adiabatic heating where the atmosphere is highly opaque in the infrared and visual (due to dust) and the surface of the planet is heated by the general circulation.

Now that the composition of the atmospheres of Mars and Venus is becoming known it is, strangely enough, earth, with only 0.3% carbon dioxide, that stands out among these inner planets as the "oddball." If, as appears likely, the inner planets were all formed the same way, they might be expected to have similar atmospheres. The explanation could be linked with the presence of life on earth; if all the carbon dioxide in our surface limestone rocks could be released into the atmosphere, its concentration would be 20 times higher than that of nitrogen now. But water trapped in limestone and other crust materials would represent 99% of the atmosphere if it had not been fixed. Any model of atmospheric formation has to account for the presence of all this water, which ought to have been lost from the top of the atmosphere during the first billion years, long before the development of life on the planet.

Ichiaque Rasool, of the Goddard Institute for Space Studies, recently suggested that earth originally lost most of its atmosphere; the water, carbon dioxide and nitrogen subsequently released from the crust by volcanic activity were reduced by free iron to hydrogen, methane and ammonia. Oxygen was retained as iron oxide. Then the presence of ammonia in the upper atmosphere shielded water molecules



How do you measure the Differential Linearity of your Pulse Height Analyzer?



It's easy with the Berkeley Nucleonics GL-3 Pulse Generator! The Model GL-3 measures differential linearity, integral linearity, resolution, and stability in pulse height analyzers and amplifiers. Two modes of operation are provided: The Sliding Pulse Mode and the Precision Pulse Mode.

Sliding Pulse Mode: In this mode (see diagram), an operational amplifier generates a ramp waveform which is sampled by a solid-state chopper. The chopper's output is a linearly increasing and decreasing sliding pulse train with a differential linearity better than 0.25%.

A test of the differential linearity of a pulse height analyzer is made by connecting the sliding pulse train to the input of the analyzer. If the analyzer is linear, an equal number of counts will be stored in each channel and the display will be a horizontal straight line. However, if there is non-linearity in the analyzer, the display will be a curved line.

Precision Pulse Mode: In this mode, the solid state chopper is connected to an adjustable dc reference voltage. The output pulses are uniform in amplitude with a temperature stability better than 0.0033°C. and an

amplitude jitter less than 0.002% rms. Amplitude is adjustable with an integral linearity of better than 0.1%.

In both the sliding pulse and precision pulse modes, the pulse frequency is adjustable at a random rate from 1 to 10 kc or a periodic rate from 1 cps to 1 mc.

For complete technical data and prices please write to Dept. 70.



Berkeley Nucleonics Corp.
1198 Tenth St., Berkeley, Calif. 94710

in lower regions from the ultraviolet radiation that dissociates them. If Mars and Venus have no iron in their crusts this process would not have happened, and water would have been lost as hydrogen atoms and hydroxyl molecules; however the presence of iron in both planets is generally assumed because of their densities.

An alternative suggestion by A. G. W. Cameron (Yeshiva University) is that earth's magnetic field made the difference; ionized hydrogen and hydroxyl radicals in the ionosphere

would have been trapped by interaction with this field. Both the American and Russian probes showed that Venus has no magnetic field, so the solar wind would easily have carried the hydrogen away. —JTS

Radio Telescopes Linked in Southern Hemisphere

Another long baseline radio interferometer is in operation, the first one in the southern hemisphere. The linkup (similar to those described in *PHYSICS TODAY*, November 1967, page 69) is between two NASA Deep Space Sta-

tions in Australia, one near Woomera in South Australia and one near Canberra in New South Wales. Its baseline is more than 9×10^6 wavelengths long at 2295 MHz. Preliminary observations of radio source 3C 273 show that the interferometer has an angular resolution better than 0.008 sec of arc.

The method by which these two telescopes are paired is applicable to any of NASA's Deep Space Stations, so that there is now the possibility of improving resolution still further by the use of pairs of instruments on different continents.

Weston Design: 400-500 GeV From a 200-GeV Accelerator

A bold new design for the 200-GeV accelerator is taking shape at the National Accelerator Laboratory. The strong-focusing synchrotron will be capable of producing 400-500 GeV protons later by addition of power supplies to boost the magnetic field.

The design is not simply a scale-up of the 33-GeV Brookhaven AGS and 28-GeV CERN PS. The beam aperture, magnets and tunnel have all been made surprisingly small; the result is a relatively inexpensive machine with great flexibility and high intensity. It is the first synchrotron to use separated-function magnets and the first to have only an extracted proton beam.

We recently visited the National Accelerator Laboratory (NAL) office in Oak Brook, Ill. and discussed Weston plans with Robert R. Wilson, director, and Edwin (Ned) Goldwasser, deputy director. Unlike typical surroundings for physicists, the office occupies the 10th floor of a Park Avenue-style building. Until recently it resembled an aircraft company, with no partitions and lots of shouting across the room. There are now separate offices, but there is still plenty of shirt-sleeved blackboard talk. And in the middle of it all is a mockup of the accelerator tunnel (3.1 meters wide by 2.4 meters high).

Congress recently approved design money of \$7.3 million.

The laboratory submitted a proposal on 15 Oct., and Wilson hopes that



SUPPORT FACILITIES for 200-GeV accelerator to be built in Weston, Ill.

construction of the \$260-million accelerator will be in the president's budget for FY 1969. If it is, the accelerator could be ready on 1 July 1972 (although Goldwasser feels this should be postponed until the 4th of July).

Accelerator. To allow for an increase in energy at a later date, the accelerator will have a rather large radius—1 km. The magnets are capable of reaching 200 GeV at 9 kG. Later, by increasing the power supply, adding more cooling and perhaps some more rf, the magnets should easily reach 18 kG to make 400-GeV protons and may be able to reach 22.5 kG to yield 500 GeV.

Accelerator experts are impressed with the compactness of the magnets and the tunnel volume to contain the magnets. The 70-GeV accelerator just completed at Serpukhov, for example, has a tunnel with four times the vol-

ume. Since magnets and tunnel are the majors costs in building a synchrotron, the Weston design gives a considerable cost saving.

Protons will be accelerated in three stages. A 200-MeV linac (very similar to the one being built to boost AGS intensity) will inject into a 10-GeV booster synchrotron with a 0.075-km radius. The booster will be a rapid-cycling injector. Every time magnetic field intensity peaks, the beam is extracted into the main ring; filling is timed just right so that at each pulse 1/13 of the ring is filled. Once filling is finished the magnetic field in the main ring is raised, and the particles are accelerated to full energy. Although a slow booster would be cheaper and allow for higher intensity, it is not yet well understood (intensity may be limited by space charge). There is still a year to change to a slow