sponse of viewers. Films (16 mm) can be obtained by title from State University Film Service, Attention: Booking Office, 135 Western Ave., Albany, N. Y.

Available films include "Some Crucial Experiments in High-Energy Physics" (Robert Adair), "The Conceptual Impact of High-Energy Physics" (Max Dresden), "Symmetry Aspects of High-Energy Physics" (Yuval Ne'eman), "Quarks in High-Energy Physics" (Victor Weisskopf), "Political and Scientific Problems Generated by High-Energy Physics" (round-table discussion with Maurice Goldhaber, Myron Good, John Toll, Ne'eman and Weisskopf), "Laser Activated Electron Transfer in Photosynthetic Systems" (Britton Chance), "Optical Properties of Solids" (Henry Ehrenreich), "Holography" (Emmett Leith), "Non-Linear Optics" (P. Pershan), "Semiconductor Lasers" (Robert Rediker), "Porous Glasses" (R. J. Charles), "Nuclear Tracks in Meteorites" (Robert Walker).

Senior Fulbright-Hays Awards Are Available in 20 Countries

Physicists may apply for university lecturing and advanced research appointments in a score of foreign countries during the 1969–70 academic year under the Fulbright-Hays Act. Applications for lectureships should be submitted before 1 May and no later than 1 June. The deadline for research applications is 1 June.

The basic requirements for these senior awards are United States citizenship, a doctoral degree or equivalent status for research, and college or university teaching experience for lecturing appointments. Proficiency in a foreign language is required only in some cases.

These awards give a maintenance allowance in local currency to cover normal living costs for the grantee and his family while in residence abroad, as well as roundtrip travel for the grantee only. The award also includes a dollar supplement or a stipend in dollars and foreign currency for lecturers named to most non-European lands.

Application forms and details on the openings and terms of awards in particular countries are available from the Committee on International Exchange of Persons, 2101 Constitution Avenue, N.W., Washington, D.C. 20418.

The chart below tells the story. We can now offer you a line of lithium-drifted Germanium detectors from stock that will provide both greater efficiency and high resolution performance in any volume up to 40 cc.

And if that gives you an extraordinary assist, consider this. We'll be happy to demonstrate the detector of your choice at your laboratory and let you draw your own conclusions about the claims we make for it. If you like it, we'll leave it and bill you later.

To take us up on our demonstration offer or for more information, write ISOTOPES, 50 Van Buren Ave., Westwood, N.J. 07675, or call Roland Kologrivov at (201) 664-7070.

Detector Volume	Co∞ Detector Resolution	Efficiency*	Peak: Compton
20cc	< 3KeV	> 2%	10:1
30cc	< 3KeV	> 3%	10:1
40cc	< 3KeV	> 4%	10:1
*Compar	ed to 3"x 3" No	(T1) scintill	ation assembly.




ISOTOPES

A TELEDYNE COMPANY

NOW WE'VE GOT
A STOCK ANSWER FOR
ALLYOUR GERMANIUM
DETECTOR

REQUIREMENTS UP TO 40 CC VOLUME.



EXTRAORDINARY.