

professor is **Frank S. Mathews** and to associate professor **Jeffold J. Burnett**. **Joseph A. Moyzis** was named assistant professor, as of September.

Victor F. Hanson has retired after 35 years with Du Pont Co. He served as the first director of the applied-physics section and of the radiation-physics laboratory. In 1963 he was named manager of engineering-physics research.

At MIT the Everett Moore Baker Award for outstanding undergraduate teaching was given to **Harry M. Schey**.

S. S. Penner, professor of engineering and physics at the University of California, San Diego, completed his term as vice chancellor for academic affairs and will continue as director of the Institute for Pure and Applied Physical Sciences.

Joining KEV Electronics as senior physicist in the advanced development group is **Francis Harper**, formerly of Bell Telephone Laboratories.

Effective as of Sept., **Joseph W. Weinberg**, professor at Case Western Reserve, will join Syracuse University as Kenan Professor of Physics. He succeeds **William R. Fredrickson**, who will continue to teach. **K. C. Wali**, senior physicist and head of the high-energy theory group at Argonne National Laboratory, was named professor. New assistant professors are **Robert Geroch**, **Yoshiaki Ueda** and **Maurice Blackmon**.

Kenneth Young, a senior at Cal Tech was awarded the Richard P. Feynman Fellowship for graduate work in high-energy physics.

OSA Ives Medal to David Rank of Pennsylvania State

The Optical Society of America will award its Frederic Ives Medal to David Rank during its October meeting in Chicago. Rank is Evan Pugh Research Professor of Physics and head of the physics department at Pennsylvania State University.

Rank has worked on atomic and molecular spectroscopy in the ultra-violet, visible and infrared regions of the spectrum and is also known for his work on Raman and Brillouin scattering. He has also worked on geometri-

cal optics, optical instrumentation, optical shop practice and testing and the application of diffraction gratings to high-resolution spectroscopy.

In 1966 he began laser studies on nonlinear and stimulated optical phenomena and has since discovered the stimulated Rayleigh scattering, stimulated thermal Rayleigh scattering and optical mixing in stimulated Brillouin scattering. He is currently an associate editor of *The Journal of the Optical Society of America*.

Given annually, the medal was endowed by the late Herbert E. Ives in honor of his father, who was a pioneer in color photography, photoengraving and three-color process printing.

Lark-Horovitz Prize Goes To Spears for Ultrasonics

The Lark-Horovitz Prize in Physics, given by Purdue University, went to David L. Spears for his studies of intense beams of ultrasonic waves in semiconductors. The award honors Karl Lark-Horovitz and is given for exceptional ability and research by a physics graduate student.

Spears received his PhD this June and is now working at the MIT Lincoln Laboratory. Working under Ralph Bray, Spears studied how ultrasonic beams can deflect the light or greatly vary their intensity as they pass through the crystals.

ANS Special Award To Arthur G. Ward

Arthur G. Ward is the recipient of the American Nuclear Society Special Award for his fuel burn-up predictions and measurements. Ward is director of the applied-physics division, Atomic Energy Limited, Chalk River Nuclear Laboratories, and has supervised the development work for the Intense Neutron Generator.

Ward has also guided the program on fuel burn-up including determination of reactivity and isotopic composition. He derived and published the first criterion for xenon instability.

Spencer Gets Gray Medal For Charged-Particle Work

Lewis V. Spencer is the first recipient of the L.H. Gray Medal given by the International Commission on Radiation Units and Measurements. Spencer, with the Center for Radiation Research, National Bureau of Standards,

We Measure Radiant Energy...



NOW DIGITALLY MEASURE UV TO FAR IR

Model 202 Infrared Radiometer

- ☐ Stable, flat response, **black body thermopile detector** with a one cm² x one steradian total effective field of view
- ☐ Readout in **watts, watts per cm² or watts per cm² per steradian**
- ☐ Automatically ranges from 10⁻⁷ watts resolution to 10⁻¹ full scale with provision for higher range levels
- ☐ Fixed absolute measurements or variable gain
- ☐ Optional filters, shutters and irises maintained at thermopile temperature to maximize stability
- ☐ Thermally floating thermopile housing eliminates the effect of ambient temperature changes.

In Addition...

- ☐ Automatic ranging over **many orders of magnitude**
- ☐ Unique, **simplified, digital data** presentation
- ☐ **BCD 1-2-4-8** and linear analog outputs (logarithmic optional) over the entire dynamic range
- ☐ **Precalibrated**, interchangeable, plug-in sensing heads
- ☐ Remote function programming capability
- ☐ Selectable fast or slow response
- ☐ Hold range and hold reading controls
- ☐ Silicon solid-state design incorporating integrated circuits
- ☐ 0.05% range calibration resistors.



440 LOGUE AVENUE
MOUNTAIN VIEW, CA. 94040
(415) 969-9230


ACCURATE DIGITAL RADIATION MEASUREMENTS — THERMAL • INFRARED • VISIBLE • ULTRAVIOLET

IT'S NOT SHAPE ALONE THAT MAKES AIRCO °KRYOCONDUCTOR™ THE IDEAL SUPERCONDUCTOR

°Kryoconductor, Airco's multi-strand superconducting material, started with our search for the ideal superconductor.

We found an answer in niobium-titanium rod, co-reduced in copper and formed in many sizes and shapes, both single and multi-strand.

Various copper-to-superconductor ratios are available, and super-



conductor strand sizes may be specified to suit your particular needs. Strands

may be as fine as one mil.

This unique new material is the forerunner of a complete line of both high and low field superconductors, all under the trade name of Airco °Kryoconductor.

For complete information, contact: Carlton Walker, Airco °Kryoconductor, Central Research Laboratories, Murray Hill, New Jersey 07971, telephone (201) 464-2400.

AIRCO
°Kryoconductor™

and a professor at Ottawa University, Kansas, will receive the award at the XIIth International Congress of Radiology. The meeting will be held in Tokyo during 6-11 Oct.

Spencer is honored for his long-term study of the theory of charged-particle penetration. His calculation on the electron slowing-down spectrum led to

an extension of Gray's theory of cavity ionization chambers, which resulted in a cavity ionization function. He also developed a mathematical treatment of the Boltzmann equation, leading to an accurate calculation of the penetration of gamma rays in matter.

The medal was established in 1967 and honors the late Louis Harold Gray, former member and vice-chairman of the commission.

Astronomer at Kitt Peak, R. W. Michie, 38, Dies

Richard W. Michie, associate astronomer at Kitt Peak National Observatory, died 27 March after a prolonged illness. Born 1931 at Long Beach, California, he received a BS in physics at Cal Tech and a PhD in astronomy at Princeton University in 1960.

He served three years on the faculty of the University of California, Berkeley before joining the staff of the space division at Kitt Peak in 1963. He was known for his contributions to the dynamical theory of star clusters and the formation of galaxies.

W. H. Barkas Developed Nuclear-Emulsion Method

Walter H. Barkas, professor at the University of California, Riverside, died on 26 March of a heart attack at the age of 56. He was born in Portland, Oregon in 1912 and received a BS degree in 1932 and a PhD in 1936 from the University of Washington.

He was best known for his pioneering work on the development of the nuclear-emulsion technique in high-energy physics, which he applied in a series of classic measurements on pion and muon properties.

His career embraced a large range of activities, including astrophysics at the University of Washington (1933-36), beta-ray spectroscopy at Columbia University (1936-38) and fission studies at Princeton University (1938-40). During World War II, he worked for the Navy on mines and explosives, first in Washington D.C. and later as assistant director of the Applied Physics Laboratory at the University of Washington. In 1947 he was appointed head scientist at the Office of Naval Research in San Francisco.

In 1950 Barkas joined Lawrence Radiation Laboratory, where he head-

ed the group using the nuclear-emulsion technique. Contributions include the measurements of the masses, lifetimes, and decay modes of K mesons and Σ and Λ hyperons. His book, *Nuclear Research Emulsions*, was published in 1963 by Academic Press and is one of the most authoritative in its field.

He joined the University of California, Riverside in 1965, where he continued his emulsion work and built up a bubble-chamber group.

Barkas was a fellow of the American Physical Society and a member of the American Association for the Advancement of Science. He was also a member of the National Academy of Sciences Committee on Nuclear Science.

Ray Lee Edwards Was Professor for 58 Years

Ray Lee Edwards, a physics professor for 58 years, died 4 April in Claremont, Calif., after a long illness. He was 83 years old.

He received numerous degrees including a BA from Oberlin College in 1908, a BS from Ohio Wesleyan University in 1909 and an MA from Ohio State University in 1916. Edwards was also awarded a PhD by the State University of Iowa in 1925 and a DSc by Denison University in 1963.

For his contributions to teaching he received the 1945 Oersted Medal, given by the American Association of Physics Teachers. He was professor and department chairman at Park College, Missouri, and later at Miami University in Ohio, where he was also professor emeritus from 1956. The physics research library at Miami was named in his honor.


Edwards's research was in mechanics, liquid viscosities and magnetism, and he also served as associate editor of the *American Journal of Physics* during 1945-47. □

Inside this new quadrupole magnet is some very interesting engineering which results in a beam of beauty. (Sample: magnetic/geometric coincidence is within ± 0.003 inch.) Mechanical symmetry is achieved by utter simplicity of fabrication and assembly - simple, but ingenious nevertheless. Magnetic symmetry follows as surely as Planck's constant is constant.

Barrel quads are traditional and they look nicer, but they have twice as many parts and require loving adjustment. This one requires none. And the magnetic field stays beautiful over the full range of current. Minimum clear aperture 2.06 inches, max. gradient 1.0 kG/in., effective pole length 9.1 inches.

We're accelerator people, too, and we supply whatever an accelerator needs. For instance, there are 17 other quads in this magnet family. See below, too. Then just ask.

FINE FOCUSING MAGNET



HV HIGH VOLTAGE ENGINEERING

EQUIPMENT DIVISION, Burlington, Mass. 01803 □ Suppliers of accelerator accessories: scattering chambers, beam profile monitors, electrostatic steerers, NMR fluxmeters, beam line plumbing, radiation-resistant metal seals, Mossbauer cryostats and furnaces targets, complete beam-handling systems, standard and custom electromagnets, sources.