

creases in breadth at the expense of depth in our undergraduate programs. Unfortunately, our undergraduate planning is not independent of our graduate programs or overall academic views. We must remember; few ideas are as dangerous as the idea that a broad interdisciplinary overview of a problem is sufficient for its solution.

TEMPLE F. SMITH

Northern Michigan University

Marquette, Michigan

11/26/79

THE AUTHOR COMMENTS: My Guest Comment on "Focusing Applied Research on Global Problems" was not intended to dilute undergraduate programs. My illustrations should have made it clear that I was interested in graduate programs, and even there, in training "global generalists" who had demonstrated ability in traditional disciplines.

ROBERT E. MARSHAK

Virginia Polytechnic Institute

Blacksburg, Virginia

12/26/79

Dual-named particle

I find it rather odd that, five years after its discovery, the J/ψ particle should still carry two names. This is all the more strange as both discoverers live in the US. I suggest that the president of The American Physical Society toss a coin at the next APS meeting and decide the name once and for all for future generations.

MAURICE M. MIZRAHI

Center for Naval Analyses

Alexandria, Virginia


11/19/79

Corrections

January 1980, page 97—Dale E. Sayers's stated professional history was in error. He joined the faculty of North Carolina State University as an assistant professor of physics in 1976 and was promoted to associate professor three years later.

—The November 1979 meeting of the APS Fluid Dynamics Division was held in Notre Dame, Indiana.

February, page 48—in the article "Organic solids: is energy-band theory enough?" by Charles B. Duke and L. B. Schein, the curve for figure 7 should be rotated 180°; the corrected figure appears below.



Circle No. 42 on Reader Service Card

For Nuclear Physics

FAST
Photomultipliers
from **EMI**



More than 60 DIFFERENT tube types are available from GENCOM, ranging in size from 3/4" to 5" O.D. Variants with 6, 10, 12, 14, BeCu-O dynodes and S-11 or bialkali cathodes are available. All types have plano-concave windows suitable for coupling to scintillators. Most types are available with fused silica windows for Cerenkov Radiation Detection and similar applications. Rise times are typically 1.6 to 2.2ns, depending on tube type and operating voltage.

Divider chains or high current fast nuclear bases, as well as potted assemblies can be supplied to customer specifications in large or small quantities.

For catalogs, prices, or technical information, contact:



EMI GENCOM INC.

80 Express St., Plainview, NY 11803 • 516-433-5900
TWX 510-221-1889

Circle No. 41 on Reader Service Card

Neutral Density Filters

Made of
best grade
optical
glass
"dye"
en masse



Available in 2" x 2" size, in densities of 0.1, 0.2, 0.3, 0.4, 0.6, 0.6, 0.7, 0.8, 0.9, 1.0, 2.0 and 3.0. Custom made instrument box holding 12 filters \$300.00 per set.

For certain usages the Absorption Type Filters are preferred to the metallic and dielectric types. Colors are stable. May be stacked together for certain densities. Optical densities are held to exceptionally close tolerances of ± 0.050 mm in thickness with densities varying in 0.1 to 0.4 inclusive ± 0.005 D; in 0.6 to 1.0 inclusive ± 0.02 D; and 2.0 to 5.0 inclusive ± 0.08 D.

OPTICS FOR INDUSTRY

ROLYN OPTICS

300 North Rolyon Place

P.O. Box 148 • Arcadia, Calif. 91006

(213) 445-6550

(213) 447-3200

Integrated AMPLIFIER/ DISCRIMINATOR Tube Socket Assembly

- TTL COMPATIBLE OUTPUT
- COMPATIBLE WITH ALL PFR PMT CHAMBERS

- TWO VERSIONS:
ADS-001: 100 KHZ
max. count rate
ADS-120: 12 MHZ
max. count rate



ADS-001 is
compatible with
the Products for
Research Model
CT-100 Digital
Counter-Timer.

Call or write for complete
performance and price information



PRODUCTS FOR RESEARCH, INC.
88 HOLTON STREET
DANVERS, MA 01923 • (617) 774-3250

Circle No. 43 on Reader Service Card