
ONE HUNDRED YEARS OF ASTRONOMY IN AMERICA

In honor of the American Astronomical Society's centennial, we celebrate American astronomy's remarkable century of technological innovation and breathtaking, ever-expanding discovery.








▲ PIONEER ASTRONOMERS Annie Jump Cannon (left) and Henrietta Swan Leavitt. Cannon classified the spectral types of 225 000 stars to produce the *Henry Draper Catalogue* in 1918–26. Leavitt's 1907–12 study of Cepheid variable stars established the first rung of the cosmic-distance ladder.

▼ ASTRONOMY BENEFACTOR Andrew Carnegie (left) stands before the Mount Wilson 60-inch telescope (then the biggest telescope in the world) and next to George Ellery Hale, the astronomer who persuaded him to help pay for it.



▼ HENRY NORRIS RUSSELL (left), one of the 20th century's most influential astronomers, and Harlow Shapley, his first graduate student. Russell published his now-famous diagram of stellar luminosity against spectral type in 1914. That same year, Shapley deduced the Sun's location in the Galaxy.





THE VELOCITY-DISTANCE RELATION AMONG EXTRA-GALACTIC NEBULAE¹

BY EDWIN HUBBLE AND MILTON L. HUMASON

ABSTRACT

Methods of determining distances of extra-galactic nebulae are discussed, and the mean absolute magnitude is revised on the basis of (1) Shapley's revision of the zero-point of the period-luminosity curve for Cepheids, and (2) more extensive observations of stars involved in nebulae. The revised value is M (vis) = -14.9. The mean color-index of the nearer extra-galactic nebulae appears to be of the order of +1.1 mag., hence M(pg) = -13.8. A color-excess is suggested which is independent of distance but shows some relation to galactic latitude.

The velocity-distance relation is re-examined with the aid of 40 new velocities, 26 of which refer to nebulae in 8 clusters or groups. Distances of the clusters, ranging out to about 32 million parsecs, have been derived from the most frequent apparent magnitudes. The velocity displacements reduce the apparent magnitudes by amounts which become appreciable for the more distant clusters.


The new data extend out to about eighteen times the distance available in the first formulation of the velocity-distance relation, but the form of the relation remains unchanged except for the revision of the unit of distance. The relation is

 $Vel. = \frac{Dist. (parsecs)}{1790}$


and the uncertainty is estimated to be of the order of 10 per cent.





FRITZ ZWICKY (chatting with Otto Stern on the right) showed in 1937 that most of the gravitational mass of the Coma cluster of galaxies is invisible. Earlier, in 1934, he and Walter Baade had identified supernovae with the explosive transformation of normal stars into neutron stars.




◆ ROBERT HERMAN (shown here), Ralph Alpher, and George Gamow developed, in the late 1940s, the physics of what is now known as the Big Bang model of the universe, which included the prediction that a microwave blackbody background radiation should pervade the cosmos.















▼ SAUL PERLMUTTER leads the Supernova Cosmology Project, which has been gathering evidence since 1995 that the expansion of the universe is accelerating—possibly due to a nonzero cosmological constant. (Photograph of Perlmutter and supernova 1987A courtesy of Lawrence Berkeley National Laboratory.)

▲ LAUNCHED IN 1983, IRAS surveyed the sky at 12, 25, 60, and 100 μm and detected more than 350 000 cosmic sources of infrared radiation. IRAS also greatly elucidated the physics of star formation, which takes place in dense dusty clouds like Rho Ophiuchi, shown on the left. (Images courtesy of Caltech/JPL/NASA.)



▲ THE DUST RING that surrounds the star HR 4796A (blotted out in this 1998 image) could harbor young planets (Hubble Space Telescope image courtesy of Brad Smith, Glenn Schneider, and NASA.)

