

Hanbury Brown and Twiss: Important, anti-weird, beautiful

Daniel Kleppner's Reference Frame entitled "Hanbury Brown's Steamroller" (PHYSICS TODAY, August 2008, page 8) provided much insight into the correlation of photon counts from separate detectors. However, more than just demonstrating an interesting physical phenomenon, the technique was important for astronomy. In combination with radiant fluxes at Earth's surface, the angular diameters of 32 hot stars measured with the intensity interferometer near Narrabri, Australia,¹ gave empirical absolute surface fluxes in $\text{erg cm}^{-2} \text{s}^{-1} \text{Hz}^{-1}$ for the stars. With the UV flux measurements from the *Orbiting Astronomical Observatory-2* and the longer-wavelength data from the ground, the angular diameters gave the stellar surface temperatures.² Previously, astronomers depended on fitting flux distributions to imperfect model atmospheres.

References

1. R. Hanbury Brown, J. Davis, L. R. Allen, *Mon. Not. R. Astron. Soc.* **137**, 375 (1967); **167**, 121 (1974).
2. A. D. Code, J. Davis, R. C. Bless, R. Hanbury Brown, *Astrophys. J.* **203**, 417 (1976).

Donald C. Morton

(don.morton@nrc.gc.ca)

National Research Council Canada
Victoria, British Columbia

I read with interest Daniel Kleppner's lucid column on Hanbury Brown's "steamroller." Kleppner mentions an aspect of the Hanbury Brown and Twiss (HBT) effect that at first seemingly defied a quantum interpretation; he terms it "anti-weird," since in certain cases a completely classical understanding is possible. Other aspects of the Hanbury Brown physics are truly weird in the

sense that they violate a Bell inequality.

I and collaborators Peter Samuelsson and Eugene Sukhorukov have proposed an interferometry experiment with fermions in an electrical conductor. Carriers are injected into the conductor from two contacts, and the cross correlation of the electrical current is measured at two other contacts along the conductor.¹ In that arrangement, it is possible to investigate the Aharonov-Bohm effect in the fermionic HBT interferometer. Our initial discussions of such an effect date back to the early 1990s.² My collaborators and I found a geometry in which the conductance (intensity) shows no Aharonov-Bohm signature, but the current correlation (intensity correlation) is nevertheless a periodic function of the Aharonov-Bohm flux.¹

We termed our finding the two-particle Aharonov-Bohm effect. The effect was recently observed by a group headed by Moty Heiblum at the Weizmann Institute of Science. His team cooled a mesoscopic conductor to tens of millikelvin and measured its conductance and noise correlations.³ Theoretical work by Carlo Beenakker at Leiden University has shown that due to orbital quasiparticle entanglement,⁴ the two-particle effect is equivalent at zero temperature to a violation of a Bell inequality.¹ The case of nonzero temperature is more complicated and is still a subject of research.⁵ The possibility of observing truly weird quantum physics is certain to increase interest in Hanbury Brown's physics.

References

1. P. Samuelsson, E. V. Sukhorukov, M. Büttiker, *Phys. Rev. Lett.* **92**, 026805 (2004).
2. M. Büttiker, *Phys. Rev. Lett.* **68**, 843 (1992); *Physica B* **175**, 199 (1991).
3. I. Neder, N. Ofek, Y. Chung, M. Heiblum, D. Mahalu, V. Umansky, *Nature* **448**, 333 (2007).
4. C. W. J. Beenakker, in *Quantum Computers, Algorithms and Chaos*, Proceedings of the International School of Physics "E. Fermi," vol. 162, G. Casati et al., eds., IOS Press, Washington, DC (2006).
5. P. Samuelsson, I. Neder, M. Büttiker, *Phys. Rev. Lett.* (in press).

Markus Büttiker

(markus.buttiker@unige.ch)

University of Geneva
Geneva, Switzerland

I found Daniel Kleppner's Reference Frame concerning the Hanbury

Brown and Twiss (HBT) effect and its applications very interesting. In particular, Kleppner mentions recent beautiful experiments using the HBT effect to explore the physics of quantum gases. The applications have been even more important than it would appear from Kleppner's piece, if one considers the role that time-dependent intensity correlations of scattered light have played in dynamical studies of critical phenomena and, more generally, of phase transitions in fluids, liquid crystals, and soft matter. Early contributions to such studies came from George Benedek (an MIT colleague of Kleppner's) in the 1960s. Herman Cummins, working in those days at the Johns Hopkins University, also made relevant contributions. In the past 40 years, thousands of papers have described the applications of intensity-correlation spectroscopy not only to statistical physics but also to physical chemistry and biology.

Concerning spatial intensity correlations, I agree with Kleppner that until a few years ago no important applications had appeared, mainly because they simply reflect, in the far field, the geometric properties of the volume from which the light is collected. However, in addition to what Kleppner describes, recent work by Marzio Giglio and coworkers (University of Milan) on near-field scattering from colloidal particles¹ promises exciting new developments.

Reference

1. See, for instance, M. Giglio, M. Carpinetti, A. Vailati, *Phys. Rev. Lett.* **85**, 1416 (2000).

Vittorio Degiorgio

(vittorio.degiorgio@unipv.it)

University of Pavia
Pavia, Italy

Kleppner replies: I thank Donald Morton for calling to my attention the measurements of a number of hot stars in the 1970s. My comment that the Hanbury Brown and Twiss (HBT) effect was never "particularly important" for astronomy reflected Hanbury Brown's own assessment of his accomplishment. I believe that my comment is defensible if the impact of intensity interferometry on astronomy is compared with, for instance, the impact of very long baseline interferometry. Vittorio Degiorgio quite correctly points out that intensity correlations have long been a tool for studying critical phenomena. The early

Letters and opinions are encouraged and should be sent by e-mail to ptletters@aip.org (using your surname as "Subject"), or by standard mail to Letters, PHYSICS TODAY, American Center for Physics, One Physics Ellipse, College Park, MD 20740-3842. Please include your name, affiliation, mailing address, e-mail address, and daytime phone number on your attachment or letter. You can also contact us online at <http://www.physicstoday.org/pt/contactus.jsp>. We reserve the right to edit submissions.

controversy about the HBT effect centered on two-particle correlations, however, and not on intensity correlations. The goal of my essay was to point out the interesting manifestations of two-particle correlations in atomic fluids.

Daniel Kleppner
(kleppner@mit.edu)

Massachusetts Institute of Technology
Cambridge

Important groundwork in opsin evolution studies

The report by Charles Day (PHYSICS TODAY, October 2008, page 20) highlights some important aspects of the study of opsin evolution. However, too much credit is given to Shozo Yokoyama's publication and work. Many people have studied the selective pressures on the sequence of amino acids that confers the molecule's function. For example, during his graduate work in the early 1990s, Wesley Toller realized the evolutionary implications and did the early studies on opsins found in squirrel fish that inhabit different depths in the ocean. Yokoyama's work cites and builds on Toller's early

graduate work. Yet Day's report does little to acknowledge the prior work and implies that Yokoyama was the first to realize the link between opsin sequence, function, and the selective pressures imposed by the environment. My intention is not to diminish Yokoyama's work but to give readers a better picture of the breadth of the field and the sources of some key discoveries.

Christian H. Fritsen
(chris.fritsen@dri.edu)
Desert Research Institute
Reno, Nevada

Volume's effect on volume

I agree with Erica Ryherd, who stated in her Quick Study on acoustical design (PHYSICS TODAY, August 2008, page 68) that small spaces can result in high sound levels that detract from the spaces' proper use. However, Ryherd repeats the mistaken belief held by many acoustical design practitioners that the volume of a space "directly influences" the overall sound energy in the space.

Ryherd uses classical diffuse field theory as the basis for the metrics—most notably reverberation time—in her discussion. That theory says that the

sound energy in an enclosure is influenced only by the power output of the source and the total sound absorption of the enclosure. She mentions that same absorption as an independent variable in her definition of reverberation time. The volume independence occurs essentially because the absorption of sound energy takes place only at the enclosure's boundaries.

Simply put, the two ways to reduce the sound level in a space are to either turn down the volume of the source or add absorption. The room's dimensions have nothing to do with it.

Jerry Hyde
(jerry@jerryhyde.com)
St. Helena, California

Ryherd replies: The aim of my Quick Study was to provide a basic overview of music-rehearsal-room design. Generally speaking, too small a room can lead to excessive sound levels. The relationship between volume and absorption is one of the many nuances that cannot be fully addressed in two pages. That is why the article highlights the importance of having a project team include an acoustical designer who can effectively optimize absorption, room layout, and other aspects of a space.

Erica Ryherd
Georgia Institute of Technology
Atlanta ■



"Sure, it's an interesting concept, but do we really need mathematical proof that Casablanca should never have been colorized?"

Q:

what do you call a
**sputter cleaning
ion source
system**

priced as low as
\$5,995?

A:

the **IG2.**

simple. reliable. inexpensive.



innovations for surface science

www.rbdinstruments.com 541.330.0723