

Also shown is the NMR protein structure, with the locations of the unexchangeable amides shaded in red.

Pintacuda and colleagues didn't need to deuterate their sample, but they did use ^{13}C and ^{15}N labeling to get information about the positions of C and N atoms.

So their method is still limited to proteins that can be grown in laboratory cell cultures—they can't, for example, harvest a protein sample directly from a human patient. Pintacuda speculates that it could one day be possible to find a solid-state NMR structure of a protein using

no isotopic labeling at all, merely the naturally present magnetic nuclei. "But resolving the proton resonances would be much harder," he says, "because we wouldn't have the information from nearby carbons and nitrogens."

Although GB1 and AP205CP are both


PHYSICS UPDATE

These items, with supplementary material, first appeared at www.physicstoday.org.

X-RAY GHOST IMAGING

In its basic form, ghost imaging is a technique that indirectly produces a likeness of an object by combining the information from two light detectors—one that views the object and one that doesn't. By splitting a speckled laser beam so that two beams strike spatially separated locations on a CCD camera screen and by placing an object in one of the beams, as shown here, re-

searchers can calculate the intensity correlations between the two signals as the beams are scanned in synchrony and construct the object's ghost image. Demonstrated in 2002 using visible light, the technique holds promise for remote atmospheric sensing because of its insensitivity to turbulence. Two research groups have now extended ghost imaging to the hard x-ray regime. Daniele Pelliccia (RMIT University in Australia) and his colleagues passed an x-ray beam produced by the European Synchrotron source through a diffracting silicon crystal grating ori-



ented to transmit part of the undiffracted beam at a slight angle from the diffracted beam. An ultrafast camera placed 20 cm downstream of the beamsplitter detected both signals—one of which was partially blocked by the imaging target, a copper wire. Because the system worked with individual synchrotron pulses, retrieving the wire's ghost image took more than just calculating intensity correlations; the group also had to filter out low-frequency artifacts associated with mechanical vibrations induced by the synchrotron pulse rate. By contrast, Hong Yu (Chinese Academy of Sciences) and colleagues periodically moved their test object—a gold film with five slits—into and out of a synchrotron's x-ray beam to create their intensity-correlated signals—no beamsplitter or frequency filtering required. Turbulence is not a problem for x-ray imaging, but radiation exposure is. Pelliccia and colleagues speculate that one could pass a weak beam through fragile tissue but use its intensity correlations with a stronger beam that never touches the tissue to produce a low-noise microscopy image. (D. Pelliccia et al., *Phys. Rev. Lett.*, in press; H. Yu et al., *Phys. Rev. Lett.*, in press.)

—RMW

HIGH-VELOCITY CLOUD BLOWS MILKY WAY BUBBLE


Spread through the disk of the Milky Way are about 20 vast bubbles of million-degree plasma wrapped in warm shells of neutral atomic hydrogen. Blowing up a so-called supershell to its typical size of 1000 light-years requires an energy, 10^{45} J, equivalent to at least 30 supernovas. Given that massive short-lived stars cluster in groups of a few dozen, it's indeed possible that supershells are inflated by supernovas. But evidence for an alternative formation scenario has been uncovered recently by Geumsook Park of Seoul National University in Korea and her collaborators. The team used a seven-band receiver at the Arecibo radio telescope in Puerto Rico to resolve the spatial and velocity structure of a supershell known as GS040. At the very center of GS040 they found a high-velocity cloud (HVC). Such clouds contain a million or so solar masses of warm gas and fly through space at speeds 50 km/s higher than is typical for stars and other galactic components. According to previous calculations, when an HVC slams into the galactic disk, it launches a supersonic expansion of hot plasma that is energetic enough to form a supershell. That appears to be what led to the formation of GS040, which, given the shell's current size, took place 5 million years ago. Where did the HVC come from, and how did it acquire its high speed? Park and her colleagues speculate that the HVC is either a fragment of a nearby galaxy that was tidally disrupted by ours or a leftover part of the intergalactic medium from which our galaxy and its neighbors formed. (G. Park et al., *Astrophys. J. Lett.* **827**, L27, 2016.)

—CD

PLASMA DISCHARGE FOR FOOD STERILIZATION

Nonequilibrium plasmas, generated at atmospheric pressure through electric discharge, are in wide commercial use, especially for cleaning surfaces and improving surface adhesion. By producing a range of reactive chemical species, both charged and neutral, they can also inactivate or kill bacteria and other pathogens.

When bacteria colonize the surface of fresh fruits and leafy vegetables, they encase themselves in a protective layer known as a biofilm. Plasmas' ability to breach biofilms has



caught the attention of the food industry. Yet for plasmas to be deployed as food decontamination tools, their ability to do the job must be understood, controllable, and reproducible. In one step toward that goal, Dawei (David) Liu, Xinpei Lu, and colleagues at China's Huazhong University of Science and Technology and at Shanghai Jiao Tong University have now used a two-dimensional discharge model to analyze the interactions between the plasma and a biofilm on the surface of an apple. In the simulations, which spanned a region 3 mm wide by 5 mm high, the apple served as a floating electrode, and a series of